

SANI-WIPE®

No Rinse Hard, Non-Porous Surface Sanitizing Wipe



PRODUCT DESCRIPTION

A fast and effective delivery method to apply a sanitizing solution to hard, non-porous, food contact surfaces. Sani Professional™ Brand Sani-Wipe® is a ready to use, disposable non-woven cloth saturated with 175 ppm quaternary ammonium chloride. It is used to sanitize hard, non-porous food contact surfaces without requiring a rinse.

CHEMICAL COMPOSITION/PRODUCT DATA

A. Chemical Composition

1. Active Ingredients:

n-Alkyl (C ₁₂ 67%, C ₁₄ 25%, C ₁₆ 7%, C ₈ , C ₁₀ , C ₁₈ 1%)		
dimethyl benzyl ammonium chloride0.0175%		
Isopropyl alcohol		
Other Ingredients		
Total100.0000%		
Does not include the weight of the wipe.		

B. Primary Skin Irritation

Results: In accordance with EPA testing guidelines, this product is classified as Toxicity Category IV meaning mild or slight irritation.

C. Eye Irritation

A Primary Ocular Irritation test was performed on New Zealand White Rabbits. One eye of each rabbit was instilled with the test substance, while the contralateral eye remained untreated and served as control. Ocular lesions were evaluated by the method of Draize at 24 and 72 hours installation. Results: Product is classified as Toxicity Category III meaning that any irritation clears in 7 days or less.

D. Acute Oral Toxicity

An Acute Oral Toxicity was performed on Sprague-Dawley derived rats. The test material was administered orally. The animals were observed initially and once daily for 14 days.

Results: Acute Oral LD50: greater than 5g/kg body weight; Toxicity Category IV.

EPA Reg. No. 9480-7



BACTERIAL ORGANISM EFFICACY

ORGANISM: Staphylococcus aureus (ATCC #6538)

Escherichia coli (ATCC #11229) Shigella boydii (ATCC #9207)

Test Method: AOAC Use Dilution Test

5% Horse Serum as soil

Exposure time: 30 Seconds

Carriers: Glass Carriers

Incubation: 24 hours @ 37 degrees C

Results: 99.999% reduction of the microorganisms

was achieved within 60 seconds

ORGANISM: Salmonella choleraseius (ATCC #10708)

Klebsiella pneumoniae (ATCC #4352)

Test Method: AOAC Use Dilution Test

5% Horse Serum as soil

Exposure time: 5 minutes @ 20 degrees C

Carriers: Glass Carriers

Incubation: 24 hours @ 37 degrees C

Results: 99.9% reduction of the microorganism was

achieved within 5 minutes

INFORMATIONAL DATA REPORT

Sani Professional™ Brand Sani-Wipe® has been proven effective against the following organisms:

MICROORGANISM	ТҮРЕ	CHARACTERISTICS
Staphylococcus aureus	Gram positive	Highly infectious, skin lesions, boils, food poisoning, Hospital acquired infection
		100a poisoning, nospital acquirea intection
Salmonella choleraesius	Gram negative	Food containment, Gastroenteritis, Septicemia
Klebsiella pneumoniae	Gram negative	Urinary tract infections, Septicemia, diarrhea
Escherichia coli	Gram negative	Infant diarrhea, urinary tract infections
Shigella boydii	Gram negative	Bacterial infection of the bowel, gastroenteritis
		bacterial dysentery, diarrhea

CAUTIONS

• Avoid contact with eyes.



ISO 9001:2000 certified
Orangeburg, NY
saniprofessional.com